

COPD Review

Webinar for
Michigan Center for Clinical Systems
Improvement (Mi-CCSI)


October 24, 2016

Karen Meyerson, MSN, APRN, NP-C, AE-C
Director, Commercial Care Management
Priority Health

Patient Education

The goal of all patient education is to help patients take the actions needed to control their asthma or COPD.

Becoming COPD Prepared: A 4-STEP Approach



www.copd.org

STEP: A Framework for COPD Care



S Screen patients at risk.



T Test and diagnose using spirometry.



E Educate your patients about COPD.



P Provide care and support.




Screening for COPD using a validated screening tool is an important step in identifying patients who may be at risk for COPD.



Top 5 Causes of Death


COPD 3rd Leading Cause of Death after Heart Disease and Cancer



Under-diagnosis of COPD in the United States

- Approximately 24 million adults have evidence of impaired lung function indicative of COPD
 - Over 12.7 million people have been diagnosed with COPD
 - Approximately half (50%) are undiagnosed
 - Most (70%) of patients with undiagnosed COPD are <65 years of age

Percent With Undiagnosed COPD



Mannino DM, et al. *MMWR Surveill Summ.* 2002;51(1):1-13.
Mannino DM, et al. *Proc Am Thorac Soc.* 2007;4(7):502-306.
Pleis JR, et al. *Vital Health Stat.* 2006;132:1-153.

Significant Burden of COPD

- Why are approximately 50% of people with COPD undiagnosed?
 - Patients typically seek medical attention at the moderate stage of COPD
 - 81% of patients already had moderate to very severe COPD at initial spirometry-confirmed diagnosis
- Each year due to COPD, there are approximately
 - 16.3 million office visits
 - 672,000 hospitalizations
- More than 22% of Medicare patients hospitalized for COPD were readmitted within 30 days of discharge; 36% of these readmissions were due to COPD

Why is Early Diagnosis of COPD Important?

- Lung damage starts early and is progressive
- Breathlessness occurs early, increases with severity, and is underestimated by patients
- Inactivity is common but must be carefully assessed by the health care provider
- Even patients with mild disease can have exacerbations
- Earlier intervention (i.e., smoking cessation) leaves patients with more lung function

Definition of COPD

- According to the ATS/ERS COPD Guidelines, COPD is...“a **preventable** and **treatable** disease state characterized by airflow limitation that is **not fully reversible**. The airflow limitation is usually **progressive** and is associated with an **abnormal inflammatory response** of the lungs to noxious particles or gases, primarily caused by cigarette smoking. Although COPD affects the lungs, it **also produces significant systemic consequences**.”
- According to the GOLD guidelines, COPD is characterized by **chronic inflammation** throughout the airways

Definitions of Chronic Bronchitis and Emphysema

- **Chronic bronchitis** is clinically defined as chronic productive cough for 3 months in each of 2 successive years in a patient in whom other causes of productive chronic cough have been excluded
- **Emphysema** is defined as abnormal, permanent enlargement of the airspaces distal to the terminal bronchioles, accompanied by destruction of their walls, yet without obvious fibrosis

COPD Is a Multicomponent Disease¹⁻³

BRONCHOCONSTRICTION

- Tightness in the smooth muscle surrounding the airways in the lungs

INFLAMMATION

- Structural changes
- Narrowing of small airways
- Increase in inflammatory cells



STRUCTURAL CHANGES

- Permanent damage to airways and alveoli

MUCOCILIARY DYSFUNCTION

- Excessive mucus production and decreased clearance

Adapted. Illustration Copyright ©2010 Nucleus Medical Art, All rights reserved. www.nucleusinc.com

1. Global Initiative for Chronic Obstructive Lung Disease. *Global Strategy for the Diagnosis, Management and Prevention of COPD*, Global Initiative for Chronic Obstructive Lung Disease (GOLD) 2010. <http://www.goldcopd.org>. Accessed April 13, 2011.

2. Agusti AG. *Respir Med*. 2005;99:670-682.


3. Rodriguez-Roisin R. *COPD*. 2005;2:253-262.

Some Diagnostic Indicators for COPD

- Dyspnea
 - Initially during exercise
 - Progresses to occur with minimal exercise or at rest
- Chronic cough
 - Intermittent early in disease process
 - Occurring primarily in morning
 - Persists throughout day with disease progression
- Sputum production
 - Any pattern of chronic sputum production may indicate COPD
- History of exposure to risk factors
 - Tobacco smoke
 - Occupational dusts and chemicals
 - Smoke from wood-burning stoves and heating fuels

Who Is the COPD Patient?

Perception³⁻⁵



Myth: COPD is a disease of the elderly¹


Myth: COPD is a disease of men²

1. Tinkelman, et al. *Am J Manag Care*. 2003;9:767-771. 2. Chapman KR. *Clin Chest Med*. 2004; 25:331-334. 3. Rennard SI. *New Engl J Med*. 2004; 350:965-966. 4. Kleinschmidt P. COPD and emphysema. Available at <http://emedicine.medscape.com/article/807143-overview>. 5. Rennard SI. *N Engl J Med*. 2004;305:965-966. Netter illustrations, with permission from Icon Learning Systems, a division of MediMedia USA, Inc. All rights reserved.




COPD in Younger Patients and Women Is on the Rise

Reality



- Reality: COPD afflicts the working-age population.
- Reality: COPD is also a disease of women.

COPD Is Not a Man's Disease




*Hypothetical Male Patient With
COPD Symptoms*

Diagnosed as COPD by
65% of physicians

*Hypothetical Female Patient
With COPD Symptoms*

Diagnosed as COPD by
49% of physicians



**COPD symptoms in women were most
commonly misdiagnosed as asthma.**

Screening patients for COPD

- The COPD Alliance¹ recommends the utilization of a simple validated² questionnaire
- COPD Population Screener™ - download at www.COPD.org
- Persons at risk should be asked to complete the screener.

COPD Population Screener™ (COPD-PS)

This survey asks questions about you, your breathing, and what you are able to do. To complete the survey, mark an X in the box that best describes your answer for each question below.

1. During the past 4 weeks, how much of the time did you feel short of breath?

None of the time	A little of the time	Some of the time	Most of the time	All of the time
☐ 0	☐ 0	☐ 1	☐ 2	☐ 2

2. Do you ever cough up any "stuff," such as mucus or phlegm?

No, never	Only with occasional colds or chest infections	Yes, a few days a month	Yes, most days a week	Yes, every day
☐ 0	☐ 0	☐ 1	☐ 1	☐ 2

3. Please select the answer that best describes you in the past 12 months. I do less than I used to because of my breathing problems.

Strongly disagree	Disagree	Unsure	Agree	Strongly agree
☐ 0	☐ 0	☐ 0	☐ 1	☐ 2

4. Have you smoked at least 100 cigarettes in your ENTIRE LIFE?

No	Yes	Don't know
☐ 0	☐ 2	☐ 0

5. How old are you?

Age 35 to 49	Age 50 to 59	Age 60 to 69	Age 70+
☐ 0	☐ 1	☐ 2	☐ 2

How to score the survey: In the spaces below, write the number that is next to your answer for each of the questions. Add the numbers to get the total score. The total score can range from 0 to 10.

#1 + #2 + #3 + #4 + #5 = TOTAL SCORE

If your total score is 5 or more, your breathing problems may be caused by chronic obstructive pulmonary disease (COPD). COPD is often referred to as chronic bronchitis and /or emphysema and is a serious lung disease that slowly gets worse over time. While COPD cannot be cured, it is treatable.

Please share the completed survey with your clinician. The higher your score, the more likely you are to have COPD. Your clinician can help evaluate your breathing problems by performing a simple breathing test, also known as spirometry.

If your total score is between 0 and 4, and you experience problems with your breathing, please share this survey with your clinician. Your clinician can help evaluate any type of breathing problem.

The COPD Alliance advocates clinician use of this, and other, validated screeners for the early detection of COPD in at risk populations.

 **copd**
ALLIANCE

COPD Population Screener is a trademark of QualityMetric Incorporated. (10/08)
Copyright ©2008 Boehringer Ingelheim Pharmaceuticals Inc. All rights reserved.
COPD Population Screener©2007 by QualityMetric Incorporated. All rights reserved. (10/10)

SVB2007048/SA000433

¹The COPD Alliance is composed of multidisciplinary societies, i.e., AANP, AAPA, ACCP, ACOFP, ACOI

² Martinez FJ et al. COPD.2008;5(2):85-95

COPD Population Screener™ (COPD-PS)

This survey asks questions about you, your breathing, and what you are able to do.
To complete the survey, mark an X in the box that best describes your answer for each question below.

1. During the past 4 weeks, how much of the time did you feel short of breath?

None of the time	A little of the time	Some of the time	Most of the time	All of the time
☐ 0	☐ 0	☐ 1	☐ 2	☐ 2

2. Do you ever cough up any "stuff," such as mucus or phlegm?

No, never	Only with occasional colds or chest infections	Yes, a few days a month	Yes, most days a week	Yes, every day
☐ 0	☐ 0	☐ 1	☐ 1	☐ 2

3. Please select the answer that best describes you in the **past 12 months**. I do less than I used to because of my breathing problems.

Strongly disagree	Disagree	Unsure	Agree	Strongly agree
☐ 0	☐ 0	☐ 0	☐ 1	☐ 2

4. Have you smoked at least 100 cigarettes in your **ENTIRE LIFE**?

No	Yes	Don't know
☐ 0	☐ 2	☐ 0

5. How old are you?

Age 35 to 49	Age 50 to 59	Age 60 to 69	Age 70+
☐ 0	☐ 1	☐ 2	☐ 2

How to score the survey: In the spaces below, write the number that is next to your answer for each of the questions.
Add the numbers to get the total score. The total score can range from 0 to 10.

#1 + #2 + #3 + #4 + #5 = TOTAL SCORE

If your total score is 5 or more, your breathing problems may be caused by chronic obstructive pulmonary disease (COPD). COPD is often referred to as chronic bronchitis and/or emphysema and is a serious lung disease that slowly gets worse over time. While COPD cannot be cured, it is treatable.

Please share the completed survey with your clinician. The higher your score, the more likely you are to have COPD. Your clinician can help evaluate your breathing problems by performing a simple breathing test, also known as spirometry.

If your total score is between 0 and 4, and you experience problems with your breathing, please share this survey with your clinician. Your clinician can help evaluate any type of breathing problem.

The COPD Alliance advocates clinician use of this, and other, validated screeners for the early detection of COPD in at risk populations.



STEP Tools on COPD.org

- **COPD Screener**

- This validated COPD screening tool uses five questions to determine a patient's risk for COPD and the potential need for spirometry testing.
- The screener is available in English and Spanish
- Treatment room poster




Spirometry is a test that measures the amount of air a patient can breathe out and the amount of time it takes to do so. Spirometry can be administered by trained office staff in a primary care setting.

Test and
diagnose using
spirometry.



Diagnosis of COPD

- Diagnosis often does not occur until the disease has progressed significantly
 - Lack of serious symptoms and poor recognition of clinical symptoms in early phase
- COPD is confirmed by performing a lung function test: post-bronchodilator spirometry.





Recommendations From the National Lung Health Education Program

- Primary care clinicians should perform an office spirometry test for the following patients:
 - Patients ≥ 45 years old who report smoking (current smokers and those who recently quit) in order to detect COPD
 - Patients with respiratory symptoms, such as chronic cough, sputum production, wheezing, or dyspnea on exertion



Spirometry Is Essential for Diagnosing COPD

If . . .

Chronic symptoms =
cough, sputum, and/or shortness of breath

And . . .

Exposure to risk factors = tobacco, occupational irritants, and/or indoor/outdoor pollution

Then . . .

Spirometry* to confirm COPD diagnosis

- FEV1/FVC <0.70
- FEV1 determines staging

*Additional testing: chest radiograph, echocardiogram, arterial blood gas, sputum analysis, CT scan.

Global Initiative for Chronic Obstructive Disease. Global strategy for the diagnosis, management, and prevention of COPD. Updated 2011.

www.goldcopd.org


Barriers to Spirometry Use

- Overcoming barriers to spirometry use may lead to increased COPD diagnosis.
 - Uncertainty about the impact of the test on outcome¹
 - Lack of training on spirometry use^{1,2}
 - Poor education on interpreting results^{1,2}
 - Time and logistical constraints³
 - Reimbursement concerns¹

Spirometry in Primary Care

- Clinical Value
 - Confirms diagnosis and assesses severity of COPD
 - Helps to differentiate asthma from COPD
 - Helps to assess response to bronchodilator therapy
- The instrument is inexpensive, easy to maintain, and well reimbursed.
- Primary care clinicians can be trained to perform accurate interpretations.

Old Spirometer



NEW
Portable
Office
Spirometers

Predicted Normal Values

Affected by:

- Age
- Height
- Sex
- Ethnic Origin




Spirometric Diagnosis of COPD

- COPD is confirmed by post-bronchodilator $FEV_1/FVC < 0.70$
- Post-bronchodilator FEV_1 measured 10-15 minutes after 2 to 4 puffs of a short-acting bronchodilator




Flow Volume Loop



Pharmacologic Therapy for Stable COPD

- All patients who are symptomatic merit a trial of drug treatment.
- Current long-acting medications can
 - Reduce symptoms
 - Increase exercise capacity
 - Reduce number and severity of exacerbations
 - Improve health status

Smoking Abstinence: Slows Progression of COPD



It is essential that you take the time to educate your patients about COPD, including symptoms, diagnosis, triggers of an exacerbation, and how to live better with the disease.

Educate your patients about COPD.



STEP Tools on COPD.org

- ✓ Flip Chart – Developed by the AANP “A Breath of Fresh Air” enables clinicians to have a thorough dialogue with their patients about everything from adherence to nutrition.
- ✓ Video to show patients the proper use of all available inhalers.



Providing care and support to COPD patients is what clinicians strive to do at the highest levels and it is also the fourth STEP in becoming COPD Prepared.

Provide
care and
support.



Role of the Care Manager in COPD

- Care Managers may assist in identifying patients complaining of cough or shortness of breath who may be at risk for respiratory disorders
- Ask patients over 40 years of age who are current or ex-smokers the following questions:
 - Do you cough regularly?
 - Do you cough up phlegm regularly?
 - Do simple chores make you short of breath?
 - Do you wheeze when you exert yourself, or at night?
 - Do you get frequent colds that persist longer than those of other people you know?
- If patients answer yes to any of these questions, consider spirometric testing

Role of the Care Manager in COPD

- Promote education on COPD and self-management techniques
- Counsel patients on the correct use of inhalers
- Care Managers should demonstrate the use of all newly prescribed devices and observe the patient's use of each device
- Question patients about their medication adherence in a nonjudgmental manner and take steps to resolve any reported problems

Role of the Care Manager in COPD

- Care Managers may assist patients in the prevention and management of COPD and COPD exacerbations by:
 - Identifying and referring patients who may have COPD
 - Educating patients on COPD and self-management techniques
 - Counseling patients on smoking cessation
 - Recommending vaccinations
 - Educating patients on signs/symptoms of COPD exacerbation
- Care Managers are well positioned to play a key role in many aspects of COPD management

Prevent COPD Exacerbations

- Defined as an acute change in a patient's baseline dyspnea, cough, and/or sputum beyond day-to-day variability, and sufficient to warrant a change in therapy
- The prevention of exacerbations is recognized as a goal in COPD disease-state management
- Frequency of exacerbations contributes to a decline in lung function and significant worsening in quality of life

Prevent COPD Exacerbations


- Changes in the following that are beyond normal day-to-day variations, are acute in onset, and may warrant a change in regular medication:
 - Baseline dyspnea (or breathlessness)
 - Cough
 - Sputum
- Increased breathlessness that may be accompanied by the following:
 - Wheezing and chest tightness
 - Increased cough and sputum
 - Change of the color and/or tenacity of sputum
 - Fever
- Tachycardia and tachypnea, malaise, insomnia, sleepiness, fatigue, depression, and confusion may accompany COPD exacerbations
- Decrease in exercise tolerance, fever, and/or new radiologic anomalies suggestive of pulmonary disease may be additional signs of COPD exacerbation

Prevent COPD Exacerbations

- Defined as an acute change in a patient's baseline dyspnea, cough, and/or sputum beyond day-to-day variability, and sufficient to warrant a change in therapy
- Evidence supports that exacerbations are acute inflammatory events superimposed on the chronic inflammation characteristic of COPD
- In a 12-month study, 77% of patients had at least 1 exacerbation
- Frequency of exacerbations contributes to a decline in lung function and significant worsening in quality of life
- The prevention of exacerbations is recognized as a goal in COPD disease-state management

Help Patients Recognize and Treat Exacerbations

- Changes in signs and symptoms from baseline:
 - Shortness of breath, even at rest
 - More wheezing, coughing, mucus
 - Mucus looks different
 - Chest tightness
 - Irritable, fatigued, no energy
 - Fever
 - Color changes
 - Rapid breathing, heart rate



Tips for Reducing Exacerbations

- Wash hands often
- Avoid close contact with people who are ill
- Get a flu shot yearly and make sure pneumonia immunization is up to date
- Use your long-term control medications daily
- Use antibiotics quickly for infections or sinus problems
- Follow the COPD Action Plan

MY COPD ACTION PLAN

It is recommended that patients and physicians /healthcare providers complete this action plan together. This plan should be discussed at each physician visit and updated as needed.



The green, yellow and red zones show groups of symptoms of COPD. The list of symptoms is not comprehensive, and you may experience other symptoms. In the "Actions" column, your healthcare provider will recommend actions for you to take based on your symptoms by checking the appropriate boxes. Your healthcare provider may write down other actions in addition to those listed here.

Green Zone: I am doing well today	Actions
• Usual activity and exercise level • Usual amounts of cough and phlegm/mucus • Sleep well at night • Appetite is good	☐ Take daily medicines ☐ Use oxygen as prescribed ☐ Continue regular exercise/diet plan ☐ At all times avoid cigarette smoke, inhaled irritants* ☐ _____ ☐ _____

Yellow Zone: I am having a bad day or a COPD flare	Actions
• More breathless than usual • I have less energy for my daily activities • Increased or thicker phlegm/mucus • Using quick relief inhaler/nebulizer more often • Swelling of ankles more than usual • More coughing than usual • I feel like I have a "chest cold" • Poor sleep and my symptoms woke me up • My appetite is not good • My medicine is not helping	☐ Continue daily medication ☐ Use quick relief inhaler every _____ hours ☐ Start an oral corticosteroid (specify name, dose and duration) ☐ _____ ☐ Start an antibiotic (specify name, dose and duration) ☐ _____ ☐ Use oxygen as prescribed ☐ Get plenty of rest ☐ Use pursed lip breathing ☐ At all times avoid cigarette smoke, inhaled irritants* ☐ Call provider immediately if symptoms don't improve*

Red Zone: I need urgent medical care	Actions
• Severe shortness of breath even at rest • Not able to do any activity because of breathing • Not able to sleep because of breathing • Fever or shaking chills • Feeling confused or very drowsy • Chest pains • Coughing up blood	☐ Call 911 or seek medical care immediately ☐ While getting help, immediately do the following: ☐ _____ ☐ _____

*The American Lung Association recommends that the providers select this action for all patients.

The information contained in this document is for educational use only. It should not be used as a substitute for professional medical advice, diagnosis or treatment.

For more information, visit www.Lung.org or call 1-800-LUNG-USA (1-800-586-4872)


Know When to Call the Doctor

- Shortness of breath or wheezing that does not resolve after using inhaler
- Change in color, smell, amount or thickness of mucus coughed up
- New or increased ankle swelling
- Awaken with shortness of breath > once/night
- Very tired and this lasts > than one/day
- Have a fever that lasts



Know When to go to the Hospital

- Confusion, slurring of speech or sleepiness during a serious lung infection
- Loss of alertness or two or more of:
 - Increase in seriousness of symptoms, such as trouble breathing at rest
 - Struggling to use your upper chest or neck muscles to try to breathe
 - A large increase in how fast your heart is beating
 - A large increase in how fast you are breathing
- Any severe shortness of breath or chest pain or any other severe symptom



Your name:

Today's date:



How is your COPD? Take the COPD Assessment Test™ (CAT)

This questionnaire will help you and your healthcare professional measure the impact COPD (Chronic Obstructive Pulmonary Disease) is having on your wellbeing and daily life. Your answers, and test score, can be used by you and your healthcare professional to help improve the management of your COPD and get the greatest benefit from treatment.

For each item below place a mark (X) in the box that best describes you currently. Be sure to only select one response for each question.

Example: I am very happy (0) **X** (1) (2) (3) (4) (5) I am very sad

		SCORE
I never cough	(0) (1) (2) (3) (4) (5)	I cough all the time
I have no phlegm (mucus) in my chest at all	(0) (1) (2) (3) (4) (5)	My chest is completely full of phlegm (mucus)
My chest does not feel tight at all	(0) (1) (2) (3) (4) (5)	My chest feels very tight
When I walk up a hill or one flight of stairs I am not breathless	(0) (1) (2) (3) (4) (5)	When I walk up a hill or one flight of stairs I am very breathless
I am not limited doing any activities at home	(0) (1) (2) (3) (4) (5)	I am very limited doing activities at home
I am confident leaving my home despite my lung condition	(0) (1) (2) (3) (4) (5)	I am not at all confident leaving my home because of my lung condition
I sleep soundly	(0) (1) (2) (3) (4) (5)	I don't sleep soundly because of my lung condition
I have lots of energy	(0) (1) (2) (3) (4) (5)	I have no energy at all
		TOTAL SCORE

COPD Assessment Test and the CAT logo is a trade mark of the GlaxoSmithKline group of companies.
© 2009 GlaxoSmithKline group of companies. All rights reserved.
Last Updated: February 24, 2012

eFigure A. COPD Assessment Test.

Reprinted with permission from GlaxoSmithKline. COPD assessment test. <http://catonline.org/eng/oh/indexEN.htm>. Accessed August 20, 2013. Copyright © GlaxoSmithKline group of companies. All rights reserved.

COPD Assessment Test (CAT)

- I never cough...I cough all the time
- I have no phlegm (mucus) in my chest at all...My chest is completely full of phlegm (mucus)
- My chest does not feel tight at all...My chest feels very tight
- When I walk up a hill or one flight of stairs I am not breathless...I am very breathless...
- I am confident leaving my home despite my lung condition...I am not confident...
- I sleep soundly...I don't sleep soundly because of my lung condition
- I have lots of energy...I have no energy at all


* Range of CAT scores from 0–40. Higher scores denote a more severe impact of COPD on a patient's life. The difference between stable and exacerbation patients was five units. No target score represents the best achievable outcome.

Role of the Care Manager in COPD

Smoking cessation is the single most effective and cost-effective intervention to reduce the risk of developing COPD and slow its progression

- Smoking cessation has the greatest capacity to influence the natural history of COPD. Health care providers should encourage all patients who smoke to quit.
- Additional benefits of smoking cessation:
 - Lung function begins to improve in 2 weeks to 3 months after quitting
 - Coughing and shortness of breath decreases in 1-9 months after quitting

Smoking Cessation: It's never too late to benefit lung function



Loss of lung function over 11 years in the Lung Health Study.


Anthonisen NR, Connett JE, Murray RP for the Lung Health Study Research Group. Smoking and Lung Function of Lung Health Study Participants after 11 Years. *Am J Respir Crit Care Med*. 2002;166:675-679.




Brief Strategies to Help the Patient Willing to Quit Smoking

- **ASK** Systematically identify all tobacco users at every visit
- **ADVISE** Strongly urge all tobacco users to quit
- **ASSESS** Determine willingness to make a quit attempt
- **ASSIST** Aid the patient in quitting
- **ARRANGE** Schedule follow-up contact

Sustained Quitters Had Lowest Mortality




Smoking Cessation Requires Multiple Attempts (8 on Average)



Pharmacotherapy for Smoking Cessation

All Smoking Cessation Therapies Have Significant Benefit



NRT = Nicotine replacement therapy.

ACCP's Tobacco Dependence Treatment Toolkit

A complete online resource for you and your patients

AMERICAN COLLEGE OF CHEST PHYSICIANS

Tobacco Dependence Treatment Toolkit

A COMPREHENSIVE RESOURCE FOR HEALTHCARE PROVIDERS AND TOBACCO-DEPENDENCE PROFESSIONALS

Now you can help your patients stop smoking and be reimbursed, using the protocols and coding information contained in this comprehensive tool kit. The user-friendly interactive online setting is a complete resource for you and your patients, containing background educational materials and clinically relevant instruments to facilitate a highly successful, proven approach to tobacco-dependence treatment.

TREATMENT TOPICS:
Select One

INFORMATION FOR:
Select One

Coding Principles

Need Reimbursement Advice?

Although not well recognized, physicians can be reimbursed for treating their patient's tobacco dependence. This tool kit provides a general framework for the relevant coding and billing principles.

[Learn more](#)

Statistics

Is tobacco use harmful?

Tobacco dependence is a fatal disease. As many as 90% of tobacco-dependent people identify tobacco use as harmful and want to reduce or stop using it.

yes no

Video

Management of the Tobacco Dependent Patient: Introductory Video

American College of Chest Physicians



Disclaimer | Committee | Contact Us | Sponsorship | Site Map | Suggested Citation | Home

© Copyright 2009-2010 American College of Chest Physicians

Visit <http://tobaccodependence.chestnet.org>

COPD Education Topics

- Pulmonary rehabilitation
- Maintain proper nutritional status
- Conserve energy and control stress
- Control breathing
- Oxygen therapy
- Support groups – Better Breathers Club



Global Initiative for Chronic Obstructive Lung Disease





Definition of COPD

- COPD, a common preventable and treatable disease, is characterized by persistent airflow limitation that is usually progressive and associated with an enhanced chronic inflammatory response in the airways and the lung to noxious particles or gases.
- Exacerbations and comorbidities contribute to the overall severity in individual patients.



Risk Factors for COPD

Genes

Exposure to particles

- Tobacco smoke
- Occupational dusts, organic and inorganic
- Indoor air pollution from heating and cooking with biomass in poorly ventilated dwellings
- Outdoor air pollution

Lung growth and development

Gender

Age


Respiratory infections

Socioeconomic status

Asthma/Bronchial hyper-reactivity

Chronic Bronchitis

Risk Factors for COPD





Diagnosis and Assessment: Key Points

- A clinical diagnosis of COPD should be considered in any patient who has dyspnea, chronic cough or sputum production, and a history of exposure to risk factors for the disease.
- Spirometry is *required* to make the diagnosis; the presence of a post-bronchodilator $FEV_1/FVC < 0.70$ confirms the presence of persistent airflow limitation and thus of COPD.



Diagnosis and Assessment: Key Points

- The goals of COPD assessment are to determine the severity of the disease, including the severity of airflow limitation, the impact on the patient's health status, and the risk of future events.
- Comorbidities occur frequently in COPD patients, and should be actively looked for and treated appropriately if present.



Diagnosis of COPD

SYMPTOMS

shortness of breath
chronic cough
sputum

EXPOSURE TO RISK FACTORS

tobacco
occupation
indoor/outdoor pollution

SPIROMETRY: Required to establish diagnosis



Assessment of Airflow Limitation: Spirometry

- Spirometry should be performed after the administration of an adequate dose of a short-acting inhaled bronchodilator to minimize variability.
- A post-bronchodilator $FEV_1/FVC < 0.70$ confirms the presence of airflow limitation.
- Where possible, values should be compared to age-related normal values to avoid over-diagnosis of COPD in the elderly.



Global Strategy for Diagnosis, Management and Prevention of COPD

Assessment of COPD: Goals

Determine the severity of the disease, its impact on the patient's health status and the risk of future events (for example exacerbations) to guide therapy. Consider the following aspects of the disease separately:

- current level of patient's symptoms
- severity of the spirometric abnormality
- frequency of exacerbations
- presence of comorbidities.



Assessment of COPD

- Assess symptoms
- Assess degree of airflow limitation using spirometry
- Assess risk of exacerbations
- Assess comorbidities



Symptoms of COPD

The characteristic symptoms of COPD are chronic and progressive dyspnea, cough, and sputum production that can be variable from day-to-day.

Dyspnea: Progressive, persistent and characteristically worse with exercise.

Chronic cough: May be intermittent and may be unproductive.

Chronic sputum production: COPD patients commonly cough up sputum.

Assessment of COPD

■ Assess symptoms

COPD Assessment Test (CAT)

or

Clinical COPD Questionnaire (CCQ)

or

mMRC Breathlessness scale



Assessment of Symptoms

COPD Assessment Test (CAT): An 8-item measure of health status impairment in COPD (<http://catestonline.org>).

Clinical COPD Questionnaire (CCQ): Self-administered questionnaire developed to measure clinical control in patients with COPD (<http://www.ccq.nl>).



Global Strategy for Diagnosis, Management and Prevention of COPD

Assessment of Symptoms

Breathlessness Measurement using the Modified
British Medical Research Council (mMRC)
Questionnaire:

Relates well to other measures of health status and
predicts future mortality risk.

Global Strategy for Diagnosis, Management and Prevention of COPD

Modified MRC (mMRC) Questionnaire

PLEASE TICK IN THE BOX THAT APPLIES TO YOU
(ONE BOX ONLY)

mMRC Grade 0. I only get breathless with strenuous exercise.

☐

mMRC Grade 1. I get short of breath when hurrying on the level
or walking up a slight hill.

☐

mMRC Grade 2. I walk slower than people of the same age on the
level because of breathlessness, or I have to stop for breath when
walking on my own pace on the level.

☐

mMRC Grade 3. I stop for breath after walking about 100 meters or
after a few minutes on the level.

☐

mMRC Grade 4. I am too breathless to leave the house or I am
breathless when dressing or undressing.

☐



Assessment of COPD

- Assess symptoms
- Assess degree of airflow limitation

Use spirometry for grading severity according to spirometry, using four grades split at 80%, 50% and 30% of predicted value



Classification of Severity of Airflow Limitation in COPD*

In patients with $FEV_1/FVC < 0.70$:

GOLD 1: Mild $FEV_1 \geq 80\%$ predicted

GOLD 2: Moderate $50\% \leq FEV_1 < 80\%$ predicted

GOLD 3: Severe $30\% \leq FEV_1 < 50\%$ predicted

GOLD 4: Very Severe $FEV_1 < 30\%$ predicted

**Based on Post-Bronchodilator FEV_1*



Assessment of COPD

- Assess symptoms
- Assess degree of airflow limitation using spirometry
- Assess risk of exacerbations
- Assess comorbidities

Use history of exacerbations and spirometry. Two exacerbations or more within the last year or an $FEV_1 < 50\%$ of predicted value are indicators of high risk. Hospitalization for a COPD exacerbation associated with increased risk of death.



Assess Risk of Exacerbations

To assess risk of exacerbations use history of exacerbations and spirometry:

- Two or more exacerbations within the last year *or* an $FEV_1 < 50\%$ of predicted value are indicators of high risk.
- One or more hospitalizations for COPD exacerbation should be considered high risk.



Global Strategy for Diagnosis, Management and Prevention of COPD

Combined Assessment of COPD


- Assess symptoms
- Assess degree of airflow limitation using spirometry
- Assess risk of exacerbations

Combine these assessments for the purpose of improving management of COPD



Global Strategy for Diagnosis, Management and Prevention of COPD

Combined Assessment of COPD






Global Strategy for Diagnosis, Management and Prevention of COPD

Combined Assessment of COPD

Assess symptoms first



CAT < 10

CAT ≥ 10

Symptoms

mMRC 0–1

mMRC ≥ 2

Breathlessness


If CAT < 10 *or* mMRC 0-1:
Less Symptoms/breathlessness (A
or C)

If CAT ≥ 10 *or* mMRC ≥ 2:
More Symptoms/breathlessness (B
or D)



Combined Assessment of COPD

Assess risk of exacerbations next




If GOLD 3 or 4 *or* ≥ 2 exacerbations per year *or* ≥ 1 leading to hospital admission:
High Risk (C or D)


If GOLD 1 or 2 *and* only 0 or 1 exacerbations per year (not leading to hospital admission):
Low Risk (A or B)

Global Strategy for Diagnosis, Management and Prevention of COPD

Combined Assessment of COPD



Combined Assessment of COPD



*When assessing risk, choose the **highest** risk according to GOLD grade or exacerbation history. One or more hospitalizations for COPD exacerbations should be considered high risk.)*

Patient	Characteristic	Spirometric Classification	Exacerbations per year	CAT	mMRC
A	Low Risk Less Symptoms	GOLD 1-2	≤ 1	< 10	0-1
B	Low Risk More Symptoms	GOLD 1-2	≤ 1	≥ 10	≥ 2
C	High Risk Less Symptoms	GOLD 3-4	≥ 2	< 10	0-1
D	High Risk More Symptoms	GOLD 3-4	≥ 2	≥ 10	≥ 2



Global Strategy for Diagnosis, Management and Prevention of COPD

Assess COPD Comorbidities

COPD patients are at increased risk for:

- Cardiovascular diseases
- Osteoporosis
- Respiratory infections
- Anxiety and Depression
- Diabetes
- Lung cancer
- Bronchiectasis

These comorbid conditions may influence mortality and hospitalizations and should be looked for routinely, and treated appropriately.



Differential Diagnosis: COPD and Asthma

COPD

- Onset in mid-life
- Symptoms slowly progressive
- Long smoking history

ASTHMA

- Onset early in life (often childhood)
- Symptoms vary from day to day
- Symptoms worse at night/early morning
- Allergy, rhinitis, and/or eczema also present
- Family history of asthma



Global Strategy for Diagnosis, Management and Prevention of COPD

Therapeutic Options: Key Points

- Appropriate pharmacologic therapy can reduce COPD symptoms, reduce the frequency and severity of exacerbations, and improve health status and exercise tolerance.
- None of the existing medications for COPD has been shown conclusively to modify the long-term decline in lung function.
- Influenza and pneumococcal vaccination should be offered depending on local guidelines.



Therapeutic Options:

COPD Medications

Beta₂-agonists

Short-acting beta₂-agonists

Long-acting beta₂-agonists

Anticholinergics

Short-acting anticholinergics

Long-acting anticholinergics

Combination short-acting beta₂-agonists + anticholinergic in one inhaler

Combination long-acting beta₂-agonist + anticholinergic in one inhaler

Methylxanthines

Inhaled corticosteroids

Combination long-acting beta₂-agonists + corticosteroids in one inhaler

Systemic corticosteroids

Phosphodiesterase-4 inhibitors

To be continued...

Webinar #6 COPD Medications

Questions?

- Call: 616-464-4816
- Email: Karen.Meyerson@priorityhealth.com
- Visit: www.goldcopd.org
www.copdfoundation.org
www.copd.org



Mi-CCSI
Center for Clinical
Systems Improvement

Thank you for attending this webinar!

REMINDER:

To receive continuing education credit,
please complete the on-line evaluation.

After you submit your on-line evaluation you
will be redirected to a webpage where you
can download your continuing education
certificate. The passcode to access this page

is: **dyspnea**

If you have issues completing the evaluation or
downloading the certificate, please email

karen.carpenter@miccsi.org or **amy.wales@miccsi.org**